arXiv:1211.6078v1 [astro-ph.SR] 26 Nov 2012

Magnetic dipole moment of SGRs and AXPs as white dwarf pulsars

J. G. Coelho[1] and M. Malheiro[1]

[1]*Departamento de Física,*
Instituto Tecnológico de Aeronáutica,
ITA, São José dos Campos,
12228-900, SP, Brazil

(Dated: November 27, 2012)

Abstract

The Anomalous X-ray Pulsars (AXPs) and Soft Gamma-ray Repeaters (SGRs) are some of the most interesting groups of pulsars that have been intensively studied in the recent years. They are seen usually as neutron stars with super strong magnetic fields, namely $B \gtrsim 10^{14}$ G. However, in the last two years two SGRs with low magnetic fields $B \sim (10^{12} - 10^{13})$ G have been detected. Moreover, fast and very magnetic WD pulsars have also been observed in the last years. Based on these new pulsar discoveries, white dwarf pulsars have been proposed as an alternative explanation to the observational features of SGRs and AXPs. The pulsar magnetic dipole moment depending only on the momentum of inertia I, and the observational properties as the period P and its first time derivative $\dot{P}$, can help to identify the scale of I for SGRs/AXPs. We discuss here the pulsar magnetic dipole moment m of SGRs and AXPs when a model based on a massive fast rotating highly magnetized white dwarf is considered. We show that the values for m obtained for almost all SGRs and AXPs are in agreement with the observed range 10^{34}emu $\leq m \leq 10^{36}$emu of isolated and polar magnetic white dwarfs, excepted for the two SGRs with low B and the AE Aquarii white dwarf pulsar where $m \simeq 10^{33}$ emu. This supports the understanding of SGRs and AXPs as belonging to a class of very fast and magnetic massive white dwarfs, perfect in line to the recent astronomical observations of fast white dwarf pulsars.

INTRODUCTION

Recently, an alternative description of the Soft Gamma-ray Repeaters (SGRs) and Anomalous X-ray Pulsars (AXPs)[1] based on rotating highly magnetized and very massive white dwarfs (WD) has been proposed by Malheiro et al. 2012[2], following previous works of Morini et al. 1988[3] and Paczynski 1990[4]. In this new description several observational features are easy understood and well explained as a consequence of the large radius of a massive white dwarf that manifests a new scale of mass density, moment of inertia, rotational energy, and magnetic dipole momentum in comparison with the case of neutron stars:

a) The existence of stable WDs can explain, the range of long rotation periods $2 \lesssim P \lesssim 12$ seconds observed in SGRs and AXPs. In particular, the fact we do not observe any SGRs/AXPs with $P \leq 2$ s is a consequence of the low surface gravity and density of the white dwarf comparing with the neutron stars ones. Boshkayev et al. 2012[5] performed a self consistent calculation of dense and very fast rotating white dwarfs and obtained the minimum rotational periods ~ 0.3, 0.5, 0.7 and 2.2 seconds for ^{4}He,^{12}C,^{16}O, and ^{56}Fe WDs respectively.

b) The long standing puzzle of the energetic balance of the SGRs and AXPs pulsars is solved: the steady X-ray luminosity L_X observed is smaller than the loss of rotational energy of the white dwarf, $L_X < \dot{E}_{\rm rot}$, because $\dot{E}_{\rm rot}$ is $\sim 10^4$ order of magnitude larger due to the WD moment of inertia in comparison with the neutron star one. SGRs and AXPs are rotation-powered massive white dwarf pulsars in complete analogy with neutron star pulsars.

c) The large steady luminosity L_X seen of $L_X \sim 10^{35}$ erg/s, for so slow pulsars ($\Omega \sim 1$ Hz) is also understood, as a consequence of the large radius of the dense WD that produces a large magnetic dipole moment, in the range of the magnetic white dwarfs, as we will discuss in this paper.

d) The large $\dot{P}$ observed (10^{-10}–10^{-13} s/s) are also consequence of the large WD magnetic dipole moment, as well of the large radius and the momentum of inertia, and not only of the magnetic field as it is the case for neutron stars in the magnetar model. Overcritical magnetic fields of the order of $(10^{14} - 10^{15})$ G are not more needed to explain the large spin-down breaking of the SGRs and AXPs pulsars: they can be understood as highly magnetized WD with large magnetic fields $(10^8 - 10^{10})$ G. Magnetic white dwarf tends to be more massive and small[6], exactly the properties shared by SGRs and AXPs as rotation powered dense white dwarfs[2].

e) The burst activity quite frequent in this sources is a consequence of their large angular velocity, near to the break-up, and close to the rotational instability. In this critical situation, the gravity can stress the star producing glitches originating the outburst and large flares observed[7, 8]. The scale of the rotational energy delivery by these glitches is also in the agreement with the outburst and giant flares energies of $(10^{41} - 10^{46})$ erg observed, since WD rotational energy is at least 10^4 larger than of the NS due to the large momentum of inertia (see Figs.(4) and (7) of Malheiro et al. 2012[2]).

f) The small population of SGRs and AXPs observed, only 23 in more than 1800 pulsars is also understood, since these WD pulsars are quite fast and very magnetic, and as a consequence difficult to be formed[9]. Moreover, astronomical observations indicate that isolated magnetic white dwarfs with high magnetic field are only 10% of the total magnetic found[6].

More specifically, for the source SGR 0418+5729 and its interpretation as a white dwarf, a crucial result has been recently obtained by Durant et al. 2011[10], where is shown that the optical emission of this source can also be explain by a hot white dwarf. The possibility of fast white dwarfs generated by supernovae Ia has also been investigated, where the rapidly rotating WD is formed shortly after the stellar formation episode, and the delay from stellar formation to explosion is basically determined by the spin-down time of the rapidly rotating merger remnant (see Ilkov and Soker 2012[9, 11]). Furthermore, recent studies predicted that the masses of high-field magnetic WDs should be larger than the average mass of WDs (see García-Berro et al. 2012[12]). Recently, its has also been proposed that radio emission from magnetars might be powered by rotation energy, similarly to what occurs in normal and high-B radio pulsars (see N. Rea et al. 2012[13]). In our understanding the high-B radio pulsars are in fact rotation-powered neutron stars[14–17], but the SGRs/AXPs have a different nature and are rotation white dwarf pulsars, as discussed here. Other discussions about radio emissions were investigated in rotating and oscillating magnetars by focusing on the main physical processes determining the position of their death lines in the $P - \dot{P}$ diagram (see Morozova et al. 2012[18]).

In this paper, we will discuss the magnetic dipole moment m of neutron stars and magnetic white dwarfs, to stress that the values obtained of m for SGRs and AXPs as white dwarfs are in agreement with the ones of polar and isolated magnetic WDs. The recent discoveries of SGR 0418+5729 (see N. Rea et al. 2010[19]) and Swift J1822.3-1606 (see N. Rea et al. 2012[20] and Livingstone et al. 2012[21]) with low magnetic field share some properties[22] with the recent detected fast WD pulsar AE Aquarii and RXJ 0648.0-4418, and the candidate EUVE J0317-855 (see Table I), to support the WD description of SGRs and AXPs as white dwarf pulsar.

THE STANDARD MAGNETIC DIPOLE MODEL FOR ROTATION-POWERED PULSARS

The magnetic dipole moment is related to the magnetic field strength at the magnetic pole of the star B_p by,

$$\mid \overrightarrow{m} \mid = \frac{B_p R^3}{2}, \tag{1}$$

where R is the star radius. If the star magnetic dipole moment is misaligned with the spin axis by an angle α, the energy per second emitted by the rotating magnetic dipole is (see, e.g., Shapiro & Teukolsky 1983[23] and references therein),

$$\dot{E}_{\rm dip} = -\frac{2}{3c^3} \mid \ddot{m} \mid^2 = -\frac{2 \mid \overrightarrow{m} \mid^2}{3c^3} \Omega^4 \sin^2\alpha, \tag{2}$$

where $\ddot{m}$ is the second derivative of the magnetic dipole moment, $\Omega = 2\pi/P$ its rotational frequency and c the speed of light. Thus, the physical quantity that dictates the scale of the electromagnetic radiated power emitted is, together with the angular rotational frequency, the magnetic dipole moment m of the star. The fundamental physical idea of the rotation-powered pulsar is that the X-ray luminosity - produced by the dipole field - can be expressed as originated from the loss of rotational energy of the pulsar,

$$\dot{E}_{\rm rot} = -4\pi^2 I \frac{\dot{P}}{P^3}, \tag{3}$$

associated to its spin-down rate $\dot{P}$, where P is the rotational period, and I is the momentum of inertia. Assuming that all of the rotational energy lost by the the star is carried away by magnetic dipole radiation, and equaling Eqs. (2) and (3) we deduce the expression of pulsar magnetic dipole moment,

$$m = \left(\frac{3c^3 I}{8\pi^2} P\dot{P} \right)^{1/2}. \tag{4}$$

Thus, from Eq. (1) we obtain the surface magnetic field at the equator B_e as [24]

$$B_e = B_p/2 = \left(\frac{3c^3 I}{8\pi^2 R^6} P\dot{P}\right)^{1/2}, \tag{5}$$

where P and $\dot{P}$ are observed properties and the moment of inertia I and the radius R of the star model dependent quantities. Within the model commonly addressed as magnetar[25, 26], based on a canonical neutron star of $M = 1.4M_\odot$ and $R = 10$ km and then $I \sim 10^{45}$g cm^2 as the source of SGRs/AXPs, and from Eqs. (4) and (5), we obtain the magnetic dipole moment and the magnetic field of the neutron star pulsar,

$$m_{\rm NS} = 3.2 \times 10^{37} (P\dot{P})^{1/2} {\rm emu}, \tag{6}$$

and

$$B_{\rm NS} = 3.2 \times 10^{19} (P\dot{P})^{1/2} {\rm G}. \tag{7}$$

Let's described all SGRs and AXPs as white dwarfs of radius $R = 3000$ km and a mass $M = 1.4M_\odot$, as recent studies of fast and very massive white dwarfs obtained (see K. Boshkayev et al. 2012[5]). Thus, these values of mass and radius generating the momentum of inertia $I \sim 1.26 \times 10^{50}$g cm^2, will be adopt hereafter in this work as the fiducial white dwarf model parameters. Using that parameters we obtain the magnetic dipole moment and the magnetic field of the white dwarf pulsar,

$$m_{\rm WD} = 1.14 \times 10^{40} (P\dot{P})^{1/2} {\rm emu}, \tag{8}$$

and

$$B_{\rm WD} = 4.21 \times 10^{14} (P\dot{P})^{1/2} {\rm G}. \tag{9}$$

These results clearly shows that the scale of the dipole magnetic moment in WD is $\sim 10^3$ times larger than for neutron stars, exactly the factor seen in the X-ray luminosity of SGRs/AXPs when compared with L_X of slow pulsars ($P \sim$1 to 10 s) as XDINS[27] and high-B radio pulsars[14–17]. Furthermore, the surface magnetic field for WDs is $\sim 10^5$ smaller than the ones of neutron stars, eliminating all the overcritical B fields advocated in the magnetar model.

OBSERVATIONS OF SGRS WITH LOW B AND WHITE DWARF PULSARS

In the last years, new astronomical observations of two SGRs with low magnetic field challenged the magnetar model of SGRs where the large magnetic field is the source of the steady X-ray luminosity observed, and also responsible for the outbursts seen as the main characteristic of these type of pulsar. Moreover, the characteristic ages $\tau = P/2\dot{P}$ of these two magnetars SGR 0418+5729 and Swift J1822.3-1606 are $\tau \sim (10^6 - 10^7)$ years, making them older comparing with all the others SGRs and AXPs $\tau \sim (10^3 - 10^4)$ years, an age usually seen as an indication for the association of SGRs/AXPs as young neutron star produced on supernova explosions[19–21].

These two sources even old and with a low surface dipole magnetic field, are still very active, with an X-ray quiescent luminosity $L_X \sim 200\dot{E}_{\rm rot}$ (where $\dot{E}$ is calculated with a neutron star momentum of inertia). To explain this energetic unbalance in the magnetar model, the toroidal magnetic field needs to be two order of magnitude larger than the surface poloidal field. Something new in the magnetar model since both magnetic fields where always large but at the same order for the other SGRs and AXPs.

Furthermore, some sources have even been tentatively proposed as candidates for white dwarf pulsars. A specific example is AE Aquarii, the first white dwarf pulsar, very fast with a short period $P = 33.08$ s[28–30]. The rapid braking of the white dwarf and the nature of pulse hard X-ray emission detected with japanese SUZAKU space telescope under these conditions can be explained in terms of spin-powered pulsar mechanism[31–33]. Although AE Aquarii is in a binary system with orbital period ~ 9.88 hr, very likely the power due to accretion of matter is inhibited by the fast rotation of the white dwarf[34].

More recently, X-ray multimirror mission (XMM)-Newton satellite had observed a white dwarf pulsar faster than AE Aquarii. Mereghetti et al. 2009[35, 36] showed that the X-ray pulsator RX J0648.0-4418 is the most massive white dwarfs currently known with mass $M = 1.28 M_\odot$ and radius $R = 3000$ km and very fast with spin period $P = 13.2$ s. That belongs to the binary systems HD 49798/RX J0648.0-4418. The luminosity $L_X \sim 10^{32}$ erg/s in this source is produced by accretion onto the white dwarf of the helium-rich matter from the wind of the companion, as discussed by Mereghetti et al. 2009. In this work, we do not consider the accretion model, instead we describe RX J0648.0-4418 as a rotation powered white dwarf, and obtain the magnetic dipole moment and magnetic field, using the above mentioned fiducial parameters of a fast rotating magnetized white dwarf.

EUVE J0317-855, is another observed WD pulsar candidate discovered as an extreme-ultraviolet (EUV) source by the *ROSAT Wide Field Camera* and *Extreme Ultraviolet Explorer EUVE* survey [37, 38]. Barstow et al. 1995[40] obtained a period of $P \sim 725$ s, which is also a fast, massive and very magnetic white dwarf with a dipole magnetic field is $B \sim 4.5 \times 10^8$ G, and a mass $(1.31 - 1.37)M_\odot$ which is relatively large compared with the typical WD mass $\sim 0.6M_\odot$. However, no pulse emission have been detected from EUVE J0317-855, which may suggest that the electron-positron creation and acceleration does not occur (see Kashiyama et al. 2011[39]). EUVE J0317-855 has a white dwarf companion, but is supposed to be no interaction between them, because of their large separation ($\gtrsim 10^3$ UA).

In Table I we compare and contrast the parameters of these three fast white dwarfs with the two SGRs with low B described in the white dwarf model presented before. We see that several features of two SGRs with low magnetic field are very similar to the ones of fast and magnetic white dwarfs recently observed. They are old, characteristic ages of Myr, low quiescent X-ray luminosity $L_X \sim (10^{31} - 10^{32})$, magnetic field of $B_{\rm WD} \sim (10^7 - 10^8)$ G and magnetic dipole moments of $m_{\rm WD} \sim (10^{33} - 10^{34})$ emu. These results give evidence for the interpretation of SGRs/AXPs as being rotating white dwarf pulsars.

DIPOLE MAGNETIC MOMENT OF NEUTRON STARS AND WHITE DWARFS

The possibility that strongly magnetized white dwarfs, could behave as neutron star pulsars with a pulse emission of high-energy photons in the X-ray to γ-ray band, originate interesting and very suggestive plots done for the first time by Terada et al.[41–43]: plots of the magnetic field strength and the dipole magnetic field as a function of the period of neutron stars and white dwarfs. In this paper, we reproduce these plots including the SGRs and AXPs as neutron stars (magnetars) and white dwarfs, using for them the fiducial parameters presented before, and including the fast WDs recently discovered.

In Fig. 1 upper panel, we present the magnetic field as a function of the period. Here we see that the magnetic field B of the SGRs and AXPs as NSs or WDs are quite different ($\sim 10^5$ order of difference) as pointed out before and also in Malheiro et al. 2012[2]. The magnetic field B of the two recent SGRs discovered (blue full circle) are comparable with the ones observed for the fast white dwarfs also ploted (black full circles). The magnetic fast white dwarfs are separated in two classes: isolated (orange triangles) and polars (blue asterisks), very magnetic with $B \sim (10^7 - 10^8)$ G, and the intermediate polars (gray triangle) with weaker field $B \sim 10^5$ G (few isolated magnetic WDs also belong to that class).

In Fig. 1 down panel, the magnetic dipole moment m is presented as a function of the rotational period. To calculate from Eq. (8) the values of m for SGRs/AXPs (as white dwarfs), and also for the three fast white dwarfs of Table I, we used the radius $R = 3000$ km in agreement with Boshkayev et al. 2012[5], whereas for the others white dwarfs we adopted $R = 10^4$ km following Terada et al. 2010 (for NSs the radius used is $R = 10$ km). This figure clearly shows that because of the small WD radius of the fast and massive white dwarf comparing to normal WD, the large magnetic field of the SGRs/AXPs as white dwarfs generated a magnetic dipole moment $10^{34} \leq m \leq 10^{36}$ emu, exactly in the range of isolated and polar magnetic white dwarfs .

This plot also shows that the property of white dwarf magnetic dipole moment range to be almost independent of the rotational period, is also reproduced by the SGRs and AXPs in the white dwarf model. This is not true for neutron stars where m increases with period P. This increase (slope) of m with P (see Fig. 1 down panel) is in fact much larger for SGRs/AXPs as NSs comparing to normal X-ray pulsars, and the magnitude of m is quite large $10^{32} \leq m \leq 10^{33}$ emu a manifestation of the overcritical magnetic fields deduced in the magnetar model.

The values for $m \sim (10^{33} - 10^{34})$ emu of the two SGRs with low B, are exactly at the same order of the three white dwarf pulsars observed (see Table I), and in the lower values of the observed isolated and polar white dwarf magnetic dipole moment range[43].

The large steady X-ray emission $L_X \sim 10^{35}$ erg/s observed in the SGRs/AXPs is now well understood as a consequence of the fast white dwarf rotation ($P \sim 10$ s), since the magnetic dipole moment m is at the same scale as the one observed for the very magnetic and not so fast white dwarfs. These evidences supports the description of SGRs and AXPs as belonging to a class of very fast and magnetic massive white dwarfs perfect in line with recent astronomical observations of fast white dwarf pulsars.

Acknowledgments

The authors acknowledges the financial support of the Brazilian agency CAPES, CNPq and FAPESP (São Paulo state agency, thematic project 2007/03633-3).

---

[1] S. Mereghetti, *A&A Rev.* **15**, 225 (2008).

[2] M. Malheiro, J. A. Rueda and R. Ruffini, *Publ. Astron. Soc. Japan* **64**, 56 (2012).

[3] M. Morini, N. R. Robba, A. Smith and M. van der Klis, *ApJ* **333**, 777 (1988).

[4] B. Paczynski, *ApJL* **365**, L9 (1990).

[5] K. Boshkayev, J. A. Rueda, R. Ruffini and I. Siutsou, **astro-ph: 1204.2070v1**, (2012).

[6] C. A. Tout, D. T. Wickramasinghe, J. Liebert, L. Ferrario and J. E. Pringle, *MNRAS* **387**, 897 (2008).

[7] V. M. Kaspi, F. P. Gavriil, P. M. Woods *et al.*, *ApJ* **588**, L93 (2003).

[8] P. M. Woods, V. M. Kaspi, C. Thompson *et al.*, *ApJ* **605**, 378 (2004).

[9] M. Ilkov and N. Soker, *MNRAS* **419**, 1695 (2012).

[10] M. Durant, O. Kargaltsev and G. G. Pavlov, *ApJ* **742**, 77 (2011).

[11] M. Ilkov and N. Soker, **arXiv:1208.0953**, (2012).

[12] E. García-Berro, S. Torres, P. Lorén-Aguilar *et al.*, **arXiv:1209.2622v1**, (2012).

[13] N. Rea, J. A. Pons, D. F. Torres and R. Turolla, *ApJ* **748**, L12 (2012).

[14] C. M. Espinoza, A. G. Lyne, M. Kramer *et al.*, *ApJ* **741**, L13 (2011).

[15] M. A. Livingstone, C.-Y. Ng, V. M. Kaspi *et al.*, *ApJ* **730**, 66 (2011).

[16] F. P. Gavriil, M. E. Gonzalez, E. V. Gotthelf *et al.*, *Science* **319**, 1802 (2008).

[17] H. S. Kumar and S. Safi-Harb *et al.*, *ApJ* **678**, L43 (2008).

[18] V. S. Morozova, B. J. Ahmedov and O. Zanotti, *MNRAS* **419**, 2147 (2012).

[19] N. Rea, P. Esposito and R. Turolla *et al.*, *Science* **330**, 944 (2010).

[20] N. Rea, G. L. Israel, P. Esposito *et al.*, *ApJ* **754**, 27 (2012).

[21] M. A. Livingstone, P. Scholz, V. M. Kaspi *et al.*, *ApJ* **743**, L38 (2012).

[22] J. G. Coelho and M. Malheiro, *Int. J. Mod. Phys. Conf. Ser.* **18**, 96 (2012).

[23] S. L. Shapiro and S. A. Teukolsky, *Black holes, white dwarfs, and neutron stars: The physics of compact objects*, New York: Wiley-Interscience (1983).

[24] A. Ferrari and R. Ruffini, *Astrophys. J. Lett.* **158**, L71+ (1969).

[25] R. C. Duncan and C. Thompson, *ApJ* **392**, L9 (1992).

[26] C. Thompson and R. C. Duncan, *MNRAS* **275**, 255 (1995).

[27] F. Haberl, *Astrophysics and Space Science* **308**, 181 (2007).

[28] Y. Terada, T. Hayashi, M. Ishida *et al.*, *Publ. Astron. Soc. Japan* **60**, 387 (2008c).

[29] J. R. P. Angel, E. F. Borra and J. D. Landstreet, *ApJS* **45**, 457 (1981).

[30] L. Ferrario and D. T. Wickramasinghe, *MNRAS* **356**, 615 (2005).

[31] N. R. Ikhsanov and N. G. Beskrovnaya, **astro-ph: 0809.1169v1**, (2008).

[32] N. R. Ikhsanov, *Astron. Astrophys.* **338**, 521 (1998).

[33] N. R. Ikhsanov and P. L. Biermann, *A&A* **445**, 305 (2006).

[34] N. R. Ikhsanov and N. G. Beskrovnaya, *Astronomy Reports* **56**, 595 (2012).

[35] S. Mereghetti, A. Tiengo, P. Esposito *et al.*, *Science* **325**, 1222 (2009).

[36] S. Mereghetti, N. La Palombara, A. Tiengo *et al.*, *ApJ* **737**, 51 (2011).

[37] B. Külebi, S. Jordan, E. Nelan *et al.*, *A&A* **524**, A36 (2010).

[38] L. Ferrario, S. Vennes, D. T. Wickramasinghe *et al.*, *MNRAS* **292**, 205 (1997).

[39] K. Kashiyama, K. Ioka and N. Kawanaka, *Physical Review D* **83**, 023002 (2011).

[40] M. A. Barstow, S. Jordan, D. O'Donoghue *et al.*, *MNRAS* **277**, 971 (1995).

[41] Y. Terada, T. Hayashi, M. Ishida *et al.*, *Advances in Space Research* **41**, 512 (2008d).

[42] Y. Terada, M. Ishida, K. Mukai *et al.*, *ESA Special Publication* **622**, 521 (2008).

[43] Y. Terada and T. Dotani, **astro-ph.HE: 1006.5274v1**, (2010).

TABLE I: Similarities of SGRs with low magnetic field and white dwarf pulsars. The observational properties of five sources: SGR 0418+5729, Swift J1822.3-1606, two observed white dwarf pulsars, and the candidate EUVE J0317-855. For the SGR 0418+5729 and Swift J1822.3-1606 the parameters P, $\dot{P}$ and L_X have been taken from the McGill online catalog at www.physics.mcgill.ca/ pulsar/magnetar/main.html. The characteristic age is given by Age $= P/2\dot{P}$ and the WD magnetic moment and surface magnetic field are calculated by Eqs. (8) and (9) respectively.

	SGR 0418+5729	Swift J1822.3-1606	AE Aquarii	RXJ 0648.0-4418	EUVE J0317-855
P(s)	9.08	8.44	33.08	13.2	725
$\dot{P}$ (10^{-14})	< 0.6	8.3	5.64	< 90	-
Age (Myr)	24	1.6	9.3	0.23	-
L_X (erg/s)	$\sim 6.2 \times 10^{31}$	$\sim 4.2 \times 10^{32}$	$\sim 10^{31}$	$\sim 10^{32}$	-
$B_{\rm WD}$(G)	$< 9.83 \times 10^{7}$	3.52×10^{8}	$\sim 5 \times 10^{7}$	$< 1.45 \times 10^{9}$	$\sim 4.5 \times 10^{8}$
$m_{\rm WD}$(emu)	2.65×10^{33}	0.95×10^{34}	$\sim 1.35 \times 10^{33}$	3.48×10^{34}	1.22×10^{34}



FIG. 1: (Color online) The figure shows scatter plot between period and magnetic field strength (upper) and magnetic dipole moment (lower) of neutrons stars and white dwarfs. The red plus sign points are the known pulsars from the ATNF online pulsar database. The green cross points are X-ray pulsars. The blue full square and the pink square correspond to SGRs and AXPs as neutron stars and white dwarfs respectively. The two blue full and open circle points are the recent observed SGR 0418+5729 and Swift J1822.3-1606 considering neutron stars and white dwarfs parameters, respectively. The black full point are the three observed white dwarf pulsar candidates: the X-ray pulsator RX J0648.0-4418, AE Aquarii and EUVE J0317-855. The gray triangles are the intermediate polar and the other points are polar and isolated white dwarfs (see Terada et al. 2008d, for details of these points).